

FIRST AND ONLY 3-IN-1 TOPICAL ACNE TREATMENT FOR PATIENTS AGES 12 AND UP



CABTREO™

(clindamycin phosphate, adapalene
and benzoyl peroxide) Topical Gel
1.2%/0.15%/3.1%

One Prescription
+
One Pump
+
Once a Day

Not an actual patient.

**Scan to learn about acne
treatment with CABTREO**



What is CABTREO?

CABTREO (clindamycin phosphate, adapalene and benzoyl peroxide) Topical Gel 1.2%/0.15%/3.1% is a prescription medicine used on the skin only (topical use) to treat acne vulgaris in adults and children 12 years of age and older. Do not use CABTREO in your mouth, eyes, or vagina.

IMPORTANT SAFETY INFORMATION

Do not use CABTREO if you have had an allergic reaction to clindamycin, adapalene, benzoyl peroxide, lincomycin, or any of the ingredients in CABTREO or have Crohn's disease, ulcerative colitis, inflammation of the colon (colitis), or severe diarrhea with antibiotic use.


Please see additional Important Safety Information throughout and full Prescribing Information, Patient Information and Instructions for Use [here](#).

Do you have acne?


Talk to your doctor



Acne is a complex skin condition that affects nearly 85% of adolescents and many adults, making it a widespread concern across age groups.



Additionally, acne is not simply a cosmetic issue, but a medical condition that is made up by many different factors.



Various factors such as genetics, hormonal changes, and external factors like stress can trigger acne.

IMPORTANT SAFETY INFORMATION (CONT'D)

Before using CABTREO, tell your healthcare provider about all your medical conditions, including if you: plan to have surgery, are pregnant or plan to become pregnant, are breastfeeding or plan to breastfeed, have other skin problems, including cuts, abrasions, sunburn, or eczema; or use other skin and topical acne products that may increase the irritation of your skin when used with CABTREO.

What are the different types of acne?

Inflammatory acne

Inflammatory acne arises when an overgrowth of skin bacteria is triggered by clogged pores, leading to inflammation. This presents as small, pinkish bumps called papules and pus-filled lesions, known as pustules. The result is red and swollen acne that can leave lasting scars even after the acne itself has subsided.



Papules



Pustules

Noninflammatory acne

Whiteheads and blackheads, often referred to as noninflammatory acne, can occur when there's an excess production of oil (sebum) in your skin which blocks your hair follicles. This creates the characteristic light and dark bumps.




Noninflammatory whiteheads



Noninflammatory blackheads

**CABTREO is used to treat acne in patients
12 years of age and older**

Please see additional Important Safety Information throughout and full Prescribing Information, Patient Information and Instructions for Use [here](#).


 **CABTREO™**
(clindamycin phosphate, adapalene
and benzoyl peroxide) Topical Gel
1.2%/0.15%/3.1%

Triple down with your treatment

Take the next step in your acne treatment journey



Not an actual patient.



With just one application a day, CABTREO is the first and only triple-combination topical gel that is formulated to treat acne in patients 12 years of age and older.

CABTREO has three key parts to help treat acne:

- Clindamycin kills certain bacteria
- Adapalene helps skin cells renew and calms redness
- Benzoyl peroxide eliminates bacteria and removes dead skin

Each ingredient offers a unique approach to managing acne.

IMPORTANT SAFETY INFORMATION (CONT'D)

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, herbal supplements, and if you take or use a medicine that contains erythromycin. CABTREO should not be used with products that contain erythromycin.

Real results for real people

In clinical studies, 50% of patients who used CABTREO had clear or almost clear skin after 12 weeks of use

UP TO

80%

LESS INFLAMMATORY ACNE

Up to 80% less inflammatory acne compared to just 56% in people who used the vehicle.

UP TO

73%

LESS NONINFLAMMATORY ACNE

Up to 73% less noninflammatory acne compared to only 49% in people who used the vehicle.

50%

WITH CLEARER SKIN

50% of the people using CABTREO ended up with clear or almost clear skin compared to 25% of people who used the vehicle.

CABTREO studies showed the majority of side effects reported by patients using the treatment as being mild or moderate.

Adverse reactions reported by >1% of CABTREO patients


More Common Side Effects with CABTREO:

- Pain where applied
- Red skin where applied
- Dry skin where applied

Less Common Side Effects with CABTREO:

- Irritated skin where applied
- Skin peeling where applied
- Rash where applied

Please see additional Important Safety Information throughout and full Prescribing Information, Patient Information and Instructions for Use [here](#).

 **CABTREO™**
(clindamycin phosphate, adapalene and benzoyl peroxide) Topical Gel
1.2%/0.15%/3.1%

See results from 12 weeks of treatment with CABTREO in clinical trials

Before-and-after photos from Phase 3 clinical trials at 12 weeks

13-year-old Asian female



Before CABTREO

After CABTREO

Individual results may vary.

IMPORTANT SAFETY INFORMATION (CONT'D)

What should I avoid while using CABTREO?

- Avoid or limit your time in sunlight, including use of sunlamps or tanning beds during treatment with CABTREO as it can make you more sensitive, and you could get severe sunburn. Use sunscreen and wear clothes including a hat that covers the treated areas of your skin if you have to be in sunlight.
- Cold weather and wind may irritate skin treated with CABTREO.
- Avoid applying CABTREO to areas with skin problems including, cuts, abrasions, sunburned skin, or eczema.
- Avoid skin products that may dry or irritate your skin.
- Avoid the use of “waxing” as a hair removal method on skin treated with CABTREO.



Not actual patients.

Before-and-after photos from Phase 3 clinical trials at 12 weeks

18-year-old Hispanic male



Before CABTREO




After CABTREO

Individual results may vary.

See more before-and-after
photos from real patients



Please see additional Important Safety
Information throughout and full Prescribing
Information, Patient Information and
Instructions for Use [here](#).


 **CABTREO™**
(clindamycin phosphate, adapalene
and benzoyl peroxide) Topical Gel
1.2%/0.15%/3.1%

It's time to take the next step in your acne treatment journey

Ask your doctor about prescription CABTREO

Discover more about the first and only 3-in-1 topical acne treatment for patients ages 12 and up.

Visit **CABTREO.com** for more information and to join the CABTREO community.



Scan to learn about savings opportunities >



IMPORTANT SAFETY INFORMATION (CONT'D)

What are the possible serious and most common side effects of CABTREO?

Tell your doctor right away if you experience side effects, including:


- **Allergic reactions:** Stop using CABTREO if you have hives, rash, severe itching, swelling of your face, eyes, lips, tongue or throat, trouble breathing, throat tightness, feeling faint, dizzy, or lightheaded.
- **Inflammation of the colon (colitis):** Stop using CABTREO if you have severe stomach (abdominal) cramps, watery diarrhea, or bloody diarrhea during treatment, and within several weeks after treatment with CABTREO.
- **Sensitivity to sunlight.** See “What should I avoid while using CABTREO”.
- **Skin irritation** at the application site is common with CABTREO and may include redness, scaling, dryness, stinging, burning, itching, and swelling.

These are not all the possible side effects of CABTREO. Call your doctor for medical advice about side effects. You may report side effects to Bausch Health US, LLC at 1-800-321-4576 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

Please see additional Important Safety Information throughout and full Prescribing Information, Patient Information and Instructions for Use [here](#).

Ortho | Dermatologics

CABTREO is a trademark of Ortho Dermatologics' affiliated entities. ©2024 Ortho Dermatologics' affiliated entities. CAB.0060.USA.23

 **CABTREO™**
(clindamycin phosphate, adapalene and benzoyl peroxide) Topical Gel
1.2%/0.15%/3.1%